



Regulatory Considerations for Zika Vaccines

Wellington Sun, M.D.

Director, Division of Vaccines and Related Product Applications
Office of Vaccine Research and Review
CBER/U.S. FDA

Zika virus in the Americas: an HHS expert consultation to
accelerate the development of countermeasures
March 28-20, 2016

Outline


- Licensure of vaccines in the U.S.
- Lessons from other flavivirus vaccines
- Specific considerations in Zika vaccine development
- Summary and pathway forward

Biologics Licensure

Section 351 of the Public Health Service Act, 42 USC 262

- Must demonstrate
 - product is safe, pure and potent; and
 - facility in which the biologic product is manufactured, processed, packed, or held meets standards designed to assure that the biologic product continues to be safe, pure and potent; ...
- Only those biologics that are demonstrated to be safe and effective, and that can be manufactured in a consistent manner will be licensed by the FDA

Development of Preventive Vaccines



Pathways for Licensure of Vaccines

- **“Traditional”**
- **Accelerated**
- **Animal Rule**

Demonstration of clinical safety required for all pathways

Approaches to demonstration of effectiveness differs among pathways

Safety Database Considerations

- Nature of the product
- Intended use
- Severity of the disease to be prevented
- Pre-clinical safety

Demonstrating Vaccine Effectiveness

“**Traditional**”- expectation of adequate and well-controlled clinical studies demonstrating:

- prevention of disease **or**
- Immunologic response, in some cases
 - scientifically well-established immunologic marker to predict protection that can be reliably measured in a validated assay
 - facilitated by an understanding of disease pathogenesis and mechanism by which vaccine prevents disease

Demonstrating Vaccine Effectiveness (cont'd)

- **Accelerated** approval – for product addressing serious or life-threatening illnesses and provide meaningful therapeutic benefit over existing therapy; adequate and well-controlled clinical trial using **surrogate** endpoint reasonably likely to predict clinical benefit; need to **verify** clinical benefit post-approval (*21 CFR Subpart E, 601.41*)
- **Animal Rule** approval – for serious or life-threatening conditions; definitive human efficacy studies not ethical or feasible; efficacy based on adequate and well-controlled animal studies; need to **verify** clinical benefit during exigency (*21 CFR Subpart H, 601.90-95*)

Immune “Correlate” and Traditional Approval

- Scientifically well-established immunologic marker to predict protection that can be reliably measured in a validated assay
- Facilitated by an understanding of disease pathogenesis and mechanism by which vaccine prevents disease

U.S. Licensed Flavivirus Vaccines

- Yellow Fever, 17D live-attenuated – 1935, later YF-Vax 1978
 - protection first demonstrated in monkey challenge model
 - safety and immunogenicity evaluated first in immune then YF-naïve subjects
 - early field studies in Brazil
 - \log_{10} neutralization index of ≥ 0.7 considered a threshold for protection based on monkey challenge

- Japanese Encephalitis, inactivated (JE-Vax) 1992, Ixiaro 2009
 - JE-Vax approval based on Thai efficacy trial in children 1-14 yo
 - Ixiaro approved based on immunogenicity (PRNT₅₀) non-inferiority to JE-Vax
 - PRNT₅₀ titer of 1:10 considered a threshold for protection based on mouse and human data.

Product development

“Begin with the end in mind”

- the Intended Use (Indication)

FULL PRESCRIBING INFORMATION: CONTENTS*

- | | | | |
|----------|--------------------------------------|-----------|--|
| 1 | INDICATIONS AND USAGE | | |
| 2 | DOSAGE AND ADMINISTRATION | | |
| 2.1 | Dosage and Schedule | 8.3 | Nursing Mothers |
| 2.2 | Administration | 8.4 | Pediatric Use |
| 2.3 | Preparation for Administration | 8.5 | Geriatric Use |
| 3 | DOSAGE FORMS AND STRENGTHS | 11 | DESCRIPTION |
| 4 | CONTRAINDICATIONS | 12 | CLINICAL PHARMACOLOGY |
| 5 | WARNINGS AND PRECAUTIONS | 12.1 | Mechanism of Action |
| 5.1 | Hypersensitivity Reactions | 13 | NONCLINICAL TOXICOLOGY |
| 5.2 | Limitations of Vaccine Effectiveness | 13.1 | Carcinogenesis, Mutagenesis, Impairment of Fertility |
| 5.3 | Altered Immunocompetence | 14 | CLINICAL STUDIES |
| 6 | ADVERSE REACTIONS | 15 | REFERENCES |
| 6.1 | Clinical Trials Experience | 16 | HOW SUPPLIED / STORAGE AND HANDLING |
| 6.2 | Post-Marketing Experience | 16.1 | How Supplied |
| 7 | DRUG INTERACTIONS | 16.2 | Storage and Handling |
| 7.1 | Use with HAVRIX | 17 | PATIENT COUNSELING INFORMATION |
| 7.2 | Use with Immunosuppressive Therapies | | |
| 8 | USE IN SPECIFIC POPULATIONS | | |
| 8.1 | Pregnancy | | |

*Sections or subsections omitted from the full prescribing information are not listed.

Zika Vaccines – Potential Target Populations

➤ Endemic Areas

- Children**
- Adolescents**
- Non-pregnant females of child-bearing age**
- Pregnant women and sexual partners**
- Others?**

➤ Non-endemic Areas

- Travelers to endemic areas**

Knowledge Gaps and Vaccine Development

- Acute Zika disease clinically resembles other co-existing arboviral diseases; definitive diagnosis requires RT-PCR during short window of low-level viremia
- Low titer of viremia suggests virus may be especially susceptible to neutralization. Can neutralizing antibody predict protection? (JEV or YF vs dengue)
- Retrospective diagnosis of Zika based on current serologic methods is difficult in those with pre-existing flaviviral antibodies due to extensive cross reactivity
- Role of NHP as infection vs disease model; other animal models?
- Duration of natural immunity and cross protection between Zika strains (serotypes?) and other flaviviruses to be defined
- Potential bi-directional modulation of Zika and dengue clinical disease, e.g. antibody-dependent enhancement (Zika < - > dengue) or cross-protection
- Incomplete characterization of risks of sequelae from infection, especially neurologic, e.g. microcephaly and Guillain-Barre syndrome

Clinical Considerations for Zika Vaccines in Pregnancy

- Unlike maternal immunizations designed to prevent neonatal or infant diseases (flu, RSV, pertussis, GBS), Zika vaccination during pregnancy is protection of the fetus *in utero*
- Major congenital malformations result from defective formation of the neural tube during the 3rd and 4th weeks
- Is sterile immunity necessary to prevent viral transmission to fetus? Protection against Zika viremia must be in place at the latest by the 3rd week, prior to neural tube formation
- Active transfer of IgG across the placenta differs by subclass and avidity. Maternal transfer of antibody very minimal before 6 weeks gestation.
- Protective neutralization antibody response typically takes 2 weeks, may be longer depending on vaccine type and regimen
- Safety and immunogenicity should be studied in pregnant females after their demonstration in non-pregnant adult females
- Need for vaccination of male partner to prevent sexual transmission during pregnancy

Dengue Vaccines in Humans 2016

- Live-attenuated (LAV)
 - Recombinant chimeras
 - Cell passaged
- Purified formalin-inactivated (PIV)
- Recombinant E-protein
- PrM/E Plasmid DNA
- Heterologous prime boost with LAV and PIV

Summary

- Vaccines are licensed based on determination of safety, purity and potency and consistency of manufacture.
- Three different approaches to demonstrating vaccine effectiveness for licensure
- Major gaps exist in knowledge in Zika virus epidemiology, diagnosis, virulence, transmission, protective immunity, interactions with other co-circulating arboviruses
- Prevention of disease due to Zika is the most direct way to demonstrating vaccine effectiveness; other approaches can be considered as knowledge accumulates
- Lessons can be drawn from experience with other successful flavivirus vaccines where licensure pathways are well-worked out
- Protecting pregnant women is a priority but demonstrating vaccine effectiveness in non-pregnant population is more direct
- Each vaccine is unique. Early discussion with FDA can expedite each specific vaccine's development.

Thank You!

U.S. FDA Office of Vaccine Research and Review

Zika Vaccine POC: *Erik Henchal, Ph.D.*
Erik.henchal@fda.hhs.gov
(301)796-1853